

RACHEL LENHART, MS<sup>1</sup> • DARRYL THELEN, PhD<sup>1,3</sup> • BRYAN HEIDERSCHEIT, PT, PhD<sup>1,3,4</sup>

# Hip Muscle Loads During Running at Various Step Rates

Distance running is a common activity for many adults; however, running-related injuries are also quite common.<sup>29,32</sup> Hip muscle weakness, in particular of the abductors and external rotators, has been implicated in a number of these

injuries, including iliotibial band syndrome<sup>9,31</sup> and patellofemoral pain,<sup>15,21,27</sup> and is frequently targeted by current prevention and treatment strategies.<sup>9,10,19,22</sup>

The biomechanical demands of the hip muscles during running have been principally inferred from joint-level analyses, such as joint moments and powers<sup>8,13,25,27</sup>;

however, the distribution to the individual muscles is more challenging to determine and largely unknown.

Hip joint moments peak during the loading response of stance phase with speeds common to those of distance running.<sup>8,25</sup> Hip extensor and abductor moments have comparable peak magnitudes (approximately 2.0 Nm·kg<sup>-1</sup>), whereas the hip rotator peak moment is considerably smaller (approximately 0.5 Nm·kg<sup>-1</sup>).<sup>13,25</sup> Determining the distribution of these joint loads to the individual muscles can be complex, as many hip muscles have moment arms about more than 1 axis<sup>20</sup> and can therefore contribute to moment production in more than 1 plane (eg, the gluteus maximus can both extend and externally rotate the hip). Further, the length of the moment arm typically changes as a function of joint position, and therefore the capacity of the muscle to contribute to a particular joint moment also varies. For example, the ability of the gluteus maximus to externally rotate the hip decreases with increased hip flexion.<sup>6</sup> Knowing the force and power produced by individual hip muscles during running may be important for understanding the biomechanical mechanisms of running-related injuries. This information could in turn enable clinicians to refine exercise selection and parameters (eg, intensity, con-

• **STUDY DESIGN:** Controlled laboratory study, cross-sectional.

• **OBJECTIVES:** To characterize hip muscle forces and powers during running, and to determine how these quantities change when altering step rate for a given running speed.

• **BACKGROUND:** Hip musculature has been implicated in a variety of running-related injuries and, as such, is often the target of rehabilitation interventions, including resistance exercises and gait retraining. The differential contributions of the hip muscles to the task of running are not well understood, and may be important for recognizing the biomechanical mechanisms of running-related injuries and refining current treatment and prevention strategies.

• **METHODS:** Thirty healthy participants ran at their preferred speed at 3 different step rates: 90%, 100%, and 110% of their preferred step rate. Whole-body kinematics and ground reaction forces were recorded. A 3-D musculoskeletal model was used to estimate muscle forces needed to produce the measured joint accelerations. Forces and powers of each muscle were compared across step-rate conditions.

• **RESULTS:** Peak force produced by the gluteus medius during running was substantially greater than that of any other hip muscle, with the majority of muscles displaying a period of negative work immediately preceding positive work. The higher running step rate led to an increase in hip flexor, hamstring, and hip extensor loading during swing, but, conversely, substantially diminished peak force and work during loading response for several hip muscles, including the gluteal muscles and piriformis.

• **CONCLUSION:** Increasing running step rate for a given running speed heightened hamstring and gluteal muscle loading in late swing, while decreasing stance-phase loading in the gluteal muscles and piriformis. These results may enable clinicians to support and refine current treatment strategies, including exercise prescription and gait retraining for running-related injuries. *J Orthop Sports Phys Ther* 2014;44(10):766-774. Epub 25 August 2014. doi:10.2519/jospt.2014.5575

• **KEY WORDS:** gait retraining, hip injury, modeling, running injury, step rate

<sup>1</sup>Department of Biomedical Engineering, University of Wisconsin-Madison, Madison, WI. <sup>2</sup>Department of Mechanical Engineering, University of Wisconsin-Madison, Madison, WI. <sup>3</sup>Department of Orthopedics and Rehabilitation, University of Wisconsin-Madison, Madison, WI. <sup>4</sup>Badger Athletic Performance, University of Wisconsin-Madison, Madison, WI. This work was supported by the University of Wisconsin Sports Medicine Classic, the National Institutes of Health (1UL2RR025012, UL1TR000427, and T32GM008692), and the National Science Foundation (0966535). The experimental protocol was approved by the University of Wisconsin-Madison's Health Sciences Institutional Review Board. The authors certify that they have no affiliations with or financial involvement in any organization or entity with a direct financial interest in the subject matter or materials discussed in the article. Address correspondence to Dr Bryan Heiderscheit, Department of Orthopedics and Rehabilitation, University of Wisconsin-Madison, 1300 University Avenue, MSC 4120, Madison, WI 53705-1532. E-mail: heiderscheit@ortho.wisc.edu • Copyright ©2014 *Journal of Orthopaedic & Sports Physical Therapy*<sup>®</sup>

traction type) to better reflect the specific demands of the activity.

Increasing running step rate for a given running speed, which conversely reduces step length, has been advocated as a rehabilitation strategy to reduce hip joint loads for those with running-related injuries, thereby promoting recovery and reducing reinjury risk.<sup>13</sup> A simple 10% increase in running step rate while maintaining preferred running speed has been shown to reduce energy absorption at the hip during the loading response, with an accompanying reduction in hip abductor and internal rotator moments.<sup>13</sup> However, these potentially beneficial alterations to hip mechanics during loading response are accompanied by increased activation of the hamstring and gluteal muscles during late swing.<sup>4</sup> Further analyses are needed to understand the individual muscle loads corresponding to these joint-level biomechanical findings.

Our primary goal was to characterize hip muscle kinetics during running in a healthy adult population, providing an estimate of the individual muscle contributions throughout the stride cycle. In addition, we sought to determine how hip muscle kinetics changed with running step rate. Based on previous electromyography (EMG) and joint-level findings,<sup>4,13</sup> we hypothesized that a higher step rate would increase hamstring and gluteus maximus muscle loading during late swing but decrease loading of the gluteal muscles in early stance.

## METHODS

### Participants

**T**HIRTY HEALTHY, RECREATIONAL runners (15 men; mean  $\pm$  SD age,  $33 \pm 14$  years; mass,  $68.6 \pm 10.9$  kg; height,  $1.75 \pm 0.11$  m) were recruited for this study. All participants had a running volume that exceeded  $24.1 \text{ km}\cdot\text{wk}^{-1}$  during the preceding 3 months. Participants had no pain while running, no history of surgery to the lower limbs, and no injury of the lower limbs in the previous 3 months. The protocol for the study was

approved by the University of Wisconsin-Madison's Health Sciences Institutional Review Board, and all volunteers provided appropriate written informed consent.

### Data Acquisition

The preferred step rate for each participant was determined during a 5-minute treadmill run at his or her preferred speed. The number of right-foot strikes was counted during a 30-second period and multiplied by 4, then recomputed over a subsequent 30-second interval to ensure consistency. Each participant was then asked to run at a prescribed step rate equal to 90%, 100%, or 110% of their preferred step rate. Step rate was controlled by having participants synchronize foot-ground contact with the beat of an audible metronome. The participant's preferred speed was kept constant across the step-rate conditions, and the condition order was randomized. Data collection did not begin until the participant was able to maintain the target step rate for a minimum of 1 minute, as determined by visual inspection. Whole-body-kinematics were recorded (200 Hz) for 15 seconds during each of the running conditions using an 8-camera, passive motion-capture system (Motion Analysis Corporation, Santa Rosa, CA). Ground reaction forces and moments were simultaneously recorded (2000 Hz) using an instrumented treadmill (Bertec Corporation, Columbus, OH). A total of 40 markers were used to track the motion of the bilateral upper and lower extremities, including 21 markers placed on anatomical landmarks and 14 placed on rigid plates strapped to the thighs and shanks.<sup>13</sup> A standing calibration trial was also collected to establish joint centers, body-segment coordinate systems, segment lengths, and local positions of tracking markers. Marker position data were low-pass filtered at 12 Hz, and ground reaction forces were low-pass filtered at 50 Hz, using fifth-order, cross-validation splines.<sup>33</sup> Five right-footed strides from each condition were extracted for this analysis.

### Data Analysis

Complete details of the musculoskeletal model and computational procedures have been previously described.<sup>17</sup> In brief, a 3-D, whole-body model with 29 degrees of freedom (DOF) was scaled to each participant. The pelvis was the base segment, with 6 DOF. The hip was a 3 DOF joint modeled as a ball in socket. The tibio-femoral joint had 1 DOF, with nonsagittal rotations and translations being a function of knee flexion.<sup>1</sup> One DOF also existed at the patellofemoral joint, with patellar position and orientation determined as a function of knee flexion angle. The ankle had 1 DOF, allowing sagittal rotation. The hip joint center in the pelvis reference frame was calibrated using a hip circumduction task and a functional joint center identification routine.<sup>24</sup> Pelvis position, orientation, and joint angles were computed at each frame of running using an inverse kinematics routine that minimized the weighted sum of squared errors between the measured and model marker positions.<sup>18</sup> Generalized coordinates of the model were fit using fifth-order, cross-validation splines,<sup>33</sup> which were then numerically differentiated to obtain generalized speeds and generalized accelerations.

Lower extremity muscle forces were estimated using a musculoskeletal model that included geometric descriptions of 96 musculotendon units acting about the low back, hip, knee, and ankle joints.<sup>1</sup> Individual muscle forces ( $F_i$ ) were assumed proportional to their activation level from zero to the individual muscle's maximum isometric force ( $a_i$ ):  $F_i = a_i \times F_{i0}$ , where  $F_{i0}$  is the assumed maximum isometric force.<sup>1</sup> A muscle-force distribution algorithm, which minimized the muscle-volume weighted sum of muscle activations squared ( $\sum V_i a_i^2$ ), was used to determine muscle forces required to generate the measured accelerations of each frame of the running cycle.<sup>12</sup> The weighting factor for each muscle was taken as that muscle's volume, which was the product of the muscle's optimal fiber length and physiological cross-sectional

area. We previously showed that estimates of muscle force patterns using this model-based approach agree well with bursts and phasing of major lower extremity muscle EMG patterns over a running stride.<sup>17</sup> Muscle power was obtained by calculating the product of muscle force and velocity, with positive and negative work determined via numerical integration. Muscle forces and powers were normalized by the participant's body mass for comparison, and only those musculo-tendon units that cross the hip joint are described.<sup>20</sup>

## Statistical Analysis

Peak muscle forces within specific periods were compared across conditions using repeated-measures analysis of variance (ANOVA), with step rate as a repeated factor. For muscles with multiple peaks in force, the periods of interest were stance, early swing, and late swing. Positive and negative work performed by the hip muscles across the entire stride cycle were compared across conditions using repeated-measures ANOVA. All post hoc analyses were completed using Tukey's honestly significant difference. Statistical analyses were completed using STATISTICA Version 6.1 (StatSoft, Inc, Tulsa, OK), with a significance level of  $P < .05$ .

## RESULTS

**P**ARTICIPANTS' PREFERRED RUNNING speeds ranged from 2.4 to 3.8 m·s<sup>-1</sup> (mean ± SD, 2.81 ± 0.38 m·s<sup>-1</sup>), and preferred step rates ranged from 156 to 192 steps per minute (mean ± SD, 174 ± 9 steps per minute). During running at preferred step rate, the largest average peak hip muscle force production occurred in the gluteus medius during the loading response of stance (32 N·kg<sup>-1</sup>) (TABLE 1). The gluteus minimus and maximus, rectus femoris, and semimembranosus reached average peak forces between 8 and 23 N·kg<sup>-1</sup> during this same period. The largest average peak hip muscle force during early swing was produced by the

TABLE 1

PEAK MUSCLE FORCES DURING RUNNING AT 90%, 100%, AND 110% OF PREFERRED STEP RATE\*


Muscle/Period	90%	100%	110%
Biceps femoris long head			
Stance	3.65 ± 1.03	3.83 ± 0.98	4.07 ± 0.97 <sup>‡</sup>
Late swing	4.69 ± 0.91 <sup>†</sup>	5.19 ± 0.94	5.25 ± 0.90 <sup>‡</sup>
Semimembranosus			
Stance	7.95 ± 2.22	8.38 ± 2.04	8.95 ± 2.08 <sup>‡</sup>
Late swing	12.12 ± 2.33 <sup>†</sup>	13.54 ± 2.35	13.98 ± 2.23 <sup>‡</sup>
Gluteus maximus			
Stance	18.01 ± 3.22 <sup>†</sup>	15.80 ± 2.89	14.57 ± 2.86 <sup>†‡</sup>
Late swing	5.41 ± 1.49 <sup>†</sup>	5.92 ± 1.60	6.04 ± 1.62 <sup>‡</sup>
Gluteus medius			
Stance	34.60 ± 5.23 <sup>†</sup>	32.05 ± 4.03	28.65 ± 3.63 <sup>†‡</sup>
Late swing	5.84 ± 1.68 <sup>†</sup>	6.71 ± 2.17	7.40 ± 2.25 <sup>†‡</sup>
Gluteus minimus			
Stance	24.25 ± 4.22 <sup>†</sup>	22.97 ± 3.65	20.82 ± 3.30 <sup>†‡</sup>
Early swing	8.54 ± 2.84 <sup>†</sup>	9.76 ± 2.99	9.52 ± 2.85 <sup>‡</sup>
Late swing	3.36 ± 1.10 <sup>†</sup>	3.86 ± 1.34	4.25 ± 1.43 <sup>†‡</sup>
Tensor fascia latae			
Early swing	1.82 ± 0.42 <sup>†</sup>	1.97 ± 0.38	2.12 ± 0.45 <sup>†‡</sup>
Rectus femoris			
Stance	9.98 ± 2.94 <sup>†</sup>	8.93 ± 2.78	7.88 ± 2.84 <sup>†‡</sup>
Early swing	7.28 ± 1.59 <sup>†</sup>	8.00 ± 1.43	9.11 ± 1.79 <sup>†‡</sup>
Sartorius			
Early swing	0.38 ± 0.09 <sup>‡</sup>	0.41 ± 0.08	0.46 ± 0.10 <sup>†‡</sup>
Psoas			
Early swing	12.78 ± 3.63	13.19 ± 2.94	13.36 ± 3.17
Iliacus			
Early swing	17.63 ± 3.96 <sup>†</sup>	19.29 ± 3.53	20.43 ± 4.16 <sup>†‡</sup>
Adductor magnus			
Stance	3.01 ± 0.97 <sup>†</sup>	2.69 ± 0.89	2.61 ± 0.97 <sup>‡</sup>
Adductor brevis			
Early swing	3.36 ± 0.86 <sup>†</sup>	3.62 ± 0.75	3.76 ± 0.76 <sup>‡</sup>
Adductor longus			
Early swing	1.57 ± 0.41	1.59 ± 0.36	1.63 ± 0.37
Piriformis			
Stance	3.47 ± 1.13	2.97 ± 0.98	2.56 ± 0.82 <sup>†‡</sup>
Early swing	1.27 ± 0.42 <sup>†</sup>	1.53 ± 0.39	1.78 ± 0.37 <sup>†‡</sup>
Late swing	1.20 ± 0.61	1.29 ± 0.57	1.34 ± 0.49

\*Values are mean ± SD N·kg<sup>-1</sup>. For muscles with multiple peaks, the range queried for each peak is listed based on the period of the running cycle.  
<sup>†</sup>Different from 100% condition ( $P < .05$ ).  
<sup>‡</sup>Different from 90% condition ( $P < .05$ ).

iliacus (19 N·kg<sup>-1</sup>) and the semimembranosus (14 N·kg<sup>-1</sup>) during late swing.

Step rate had minimal effect on the temporal pattern of muscle forces (FIGURE 1) but did alter peak forces produced (TABLE 1; APPENDIX TABLE 1, available online).

During loading response, peak forces of the gluteal muscles, rectus femoris, adductor magnus, and piriformis significantly decreased as step rate increased. In contrast, the biceps femoris long head and semimembranosus peak forces in-



**FIGURE 1.** Mean muscle force (N·kg<sup>-1</sup>) production across the running cycle. The shaded area indicates 1 SD of force produced across all participants at the preferred (100%) step-rate condition. The other step rates (90% and 110% of preferred step rate) had similar variance (not shown). The vertical dashed line indicates toe-off. \*All conditions were different from one another ( $P < .05$ ). †90% different from 110% ( $P < .05$ ). ‡90% different from 100% and 110% ( $P < .05$ ). Abbreviations: AddBrev, adductor brevis; AddLong, adductor longus; AddMag, adductor magnus; BFL, biceps femoris long head; GMax, gluteus maximus; GMed, gluteus medius; GMin, gluteus minimus; RF, rectus femoris; SM, semimembranosus; TFL, tensor fascia latae.

creased with step rate during this period. During early swing, several muscles showed an increase in force when the step rate was increased, including the tensor fascia latae, gluteus minimus, rectus femoris, and adductor brevis. During late swing, the hamstrings and gluteal muscles produced higher forces as step rate increased.

Unique force and velocity profiles between the muscles led to distinctive mean power production (FIGURE 2) and work values (TABLE 2; APPENDIX TABLE 2, available online) across the running cycle. While the hamstrings performed only positive work during loading response, the glu-

teals and piriformis performed negative work for a period prior to positive work. These muscles tended to produce less net work (positive plus negative) across the running cycle as step rate increased. The psoas, iliacus, and adductor brevis performed negative work in late stance and positive work in early swing. During early swing, the positive work from the iliacus increased with step rate, whereas the negative work performed by the tensor fascia latae decreased as step rate increased. The rectus femoris performed largely negative work in loading response and early swing, with reduced work as step rate increased from the 90% condi-

tion. In late swing, the hamstring muscles performed negative work, followed by positive work just before foot-ground contact. Both the biceps femoris long head and the semimembranosus performed approximately 10% more negative work with the increase in step rate over the preferred step rate.

## DISCUSSION

**T**HE PURPOSE OF THIS STUDY WAS TO characterize hip muscle forces and powers produced during running, and to determine the effect of changing step rate. As hypothesized, increasing

TABLE 2

POSITIVE AND NEGATIVE WORK PERFORMED BY EACH MUSCLE ACROSS THE RUNNING CYCLE AT 3 DIFFERENT STEP-RATE CONDITIONS (90%, 100%, AND 110% OF PREFERRED)\*

Muscle/Work	90%	100%	110%
Biceps femoris long head			
Positive	0.044 ± 0.017	0.045 ± 0.015	0.045 ± 0.015
Negative	-0.055 ± 0.014	-0.056 ± 0.015	-0.060 ± 0.017 <sup>†‡</sup>
Semimembranosus			
Positive	0.110 ± 0.038	0.100 ± 0.034	0.100 ± 0.034
Negative	-0.170 ± 0.041 <sup>†</sup>	-0.180 ± 0.042	-0.200 ± 0.048 <sup>†‡</sup>
Gluteus maximus			
Positive	0.210 ± 0.066 <sup>†</sup>	0.170 ± 0.058	0.150 ± 0.048 <sup>‡</sup>
Negative	-0.051 ± 0.030 <sup>†</sup>	-0.033 ± 0.036	-0.022 ± 0.021 <sup>†‡</sup>
Gluteus medius			
Positive	0.260 ± 0.063 <sup>†</sup>	0.210 ± 0.048	0.170 ± 0.042 <sup>†‡</sup>
Negative	-0.110 ± 0.038 <sup>†</sup>	-0.100 ± 0.033	-0.073 ± 0.030 <sup>†‡</sup>
Gluteus minimus			
Positive	0.150 ± 0.049 <sup>†</sup>	0.130 ± 0.044	0.100 ± 0.038 <sup>†‡</sup>
Negative	-0.099 ± 0.033	-0.099 ± 0.029	-0.080 ± 0.025 <sup>†‡</sup>
Tensor fascia latae			
Positive	0.013 ± 0.005	0.013 ± 0.005	0.012 ± 0.005
Negative	-0.032 ± 0.010	-0.032 ± 0.010	-0.030 ± 0.001 <sup>†‡</sup>
Rectus femoris			
Positive	0.080 ± 0.029 <sup>†</sup>	0.074 ± 0.025	0.067 ± 0.023 <sup>†‡</sup>
Negative	-0.370 ± 0.093 <sup>†</sup>	-0.350 ± 0.088	-0.340 ± 0.082 <sup>‡</sup>
Sartorius			
Positive	0.015 ± 0.004	0.015 ± 0.003	0.016 ± 0.003
Negative	-0.005 ± 0.002	-0.004 ± 0.002	-0.003 ± 0.001 <sup>†‡</sup>
Psoas			
Positive	0.099 ± 0.031	0.100 ± 0.024	0.100 ± 0.022
Negative	-0.043 ± 0.019	-0.039 ± 0.015	-0.035 ± 0.016 <sup>‡</sup>
Iliacus			
Positive	0.150 ± 0.033 <sup>†</sup>	0.170 ± 0.035	0.190 ± 0.037 <sup>†‡</sup>
Negative	-0.089 ± 0.030 <sup>†</sup>	-0.079 ± 0.030	-0.075 ± 0.030 <sup>‡</sup>
Adductor magnus			
Positive	0.033 ± 0.014 <sup>†</sup>	0.030 ± 0.014	0.027 ± 0.014 <sup>‡</sup>
Negative	-0.007 ± 0.005	-0.006 ± 0.004	-0.004 ± 0.003 <sup>†‡</sup>
Adductor brevis			
Positive	0.035 ± 0.014	0.038 ± 0.014	0.037 ± 0.013
Negative	-0.021 ± 0.011	-0.019 ± 0.009	-0.015 ± 0.007 <sup>†‡</sup>
Adductor longus			
Positive	0.026 ± 0.009	0.028 ± 0.010	0.027 ± 0.009
Negative	-0.015 ± 0.006 <sup>†</sup>	-0.013 ± 0.005	-0.011 ± 0.004 <sup>†‡</sup>
Piriformis			
Positive	0.017 ± 0.010 <sup>†</sup>	0.013 ± 0.008	0.011 ± 0.006 <sup>†‡</sup>
Negative	-0.013 ± 0.006 <sup>†</sup>	-0.011 ± 0.005	-0.010 ± 0.004 <sup>‡</sup>

\*Values are mean ± SD J·kg<sup>-1</sup>.

<sup>†</sup>Different from 100% condition (P<.05).


<sup>‡</sup>Different from 90% condition (P<.05).

running step rate heightened hamstring and gluteus maximus muscle loading in late swing, an effect likely reflecting the greater limb decelerations needed to position the limb for foot-ground contact. However, after foot strike, an increased step rate results in a more erect limb posture,<sup>13,17</sup> which lessens the hip muscle forces and powers needed in the loading-response phase of stance. The decreased loading was particularly evident in the gluteal muscles and piriformis, which are muscles often implicated in running injuries.<sup>2,9,16,23,29</sup>

We previously found that model-estimated muscle forces were in temporal agreement with activation patterns from experimentally obtained EMG,<sup>17</sup> giving us confidence in our muscle force and power estimates. Further, the peak forces and temporal patterns of the hamstrings, gluteus maximus and medius, psoas, and iliacus were in general agreement with those reported at a similar running speed (3.5 m·s<sup>-1</sup>).<sup>7</sup>

As expected, the gluteal muscles produced peak forces during the loading response of stance, when hip extensor, abductor, and internal rotator joint moments are known to peak.<sup>25</sup> However, despite extensor and abductor moments being comparable in magnitude, peak forces between the 3 gluteal muscles were quite different. The sum of peak forces from the gluteus medius and minimus, 2 primary hip abductors,<sup>20</sup> was 3.5 times that of the gluteus maximus, a primary hip extensor. This disparity is not attributable to moment arm differences between the muscles, as the extension moment arm of the gluteus maximus is generally comparable to the abduction moment arm of the gluteus medius and minimus.<sup>20</sup> The disparity is more likely due to the gluteus medius and minimus being better aligned to generate the required triaxial hip joint moments in running. This was particularly evident for the anterior fibers of the gluteus medius, which have a larger internal rotation moment arm than the more posterior fibers and were recruited to a greater extent in





**FIGURE 2.** Mean hip muscle power ( $\text{W}\cdot\text{kg}^{-1}$ ) across the running cycle. The shaded area indicates 1 SD of power produced across all participants at the preferred (100%) step-rate condition. The other step rates (90% and 110% of preferred) had similar variance (not shown). The vertical dashed line indicates toe-off. Abbreviations: AddBrev, adductor brevis; AddLong, adductor longus; AddMag, adductor magnus; BFL, biceps femoris long head; GMax, gluteus maximus; GMed, gluteus medius; GMin, gluteus minimus; RF, rectus femoris; SM, semimembranosus; TFL, tensor fascia latae.

the loading response (**FIGURE 3**). In addition to the gluteus maximus, the adductor magnus, hamstrings, and more posterior fibers of the gluteus medius and gluteus minimus also contributed to the hip extensor moment in stance. Similar observations were made by Dorn et al,<sup>7</sup> who estimated that the gluteus maximus contributes about half of the hip extensor moment during the loading response of running at  $3.5 \text{ m}\cdot\text{s}^{-1}$ .

Step rate had a marked effect on hip muscle forces and powers in stance. The gluteal muscle forces decreased in proportion to the increase in step rate. That is, a 10% increase in step rate resulted in approximately a 10% decrease in peak

force from each of the gluteal muscles, with a corresponding reduction in negative and positive work. The gluteal muscles function to decrease forward speed of the body's center of mass during early stance, and, in combination with the adductor magnus, provide nearly half of the peak vertical support of body weight.<sup>11</sup> Because both the braking impulse and vertical displacement of the body's center of mass are reduced when running at a higher step rate,<sup>13</sup> the functional demands placed on the gluteal muscles (as well as adductor magnus) are likely reduced.


Step rate had a substantial impact on piriformis muscle force and power. Simi-

lar to the gluteal muscles, the piriformis reached peak force during the initial half of stance, with the muscle performing a period of negative work immediately preceding a period of positive work. Increasing step rate 10% above the preferred rate resulted in a 14% average reduction in peak piriformis force. Post hoc analysis of the negative work occurring only during initial stance revealed a nearly 40% reduction, indicating the eccentric load to the piriformis to be reduced by more than a third. However, there was a slight increase in negative work during early swing with an increase in step rate, which tempered the benefits of reduced negative work. Still, the overall negative work

progressively decreased with an increase in step rate. With excessive stretch and load suggested as causative factors of running-related piriformis pain or syndrome,<sup>2,16,30</sup> increasing running step rate may be a simple method of reducing the stretch and negative work performed by this muscle, thereby limiting injury risk and potentially being a mode of therapy. Future work should investigate this possibility.

Several muscles required greater force production during early swing when step rate was higher, including the iliacus, rectus femoris, sartorius, tensor fascia latae, gluteus minimus, and adductor brevis. This finding is somewhat expected, as each of these muscles is considered a hip flexor either as a primary or secondary action,<sup>20</sup> and each is suggested to have a key role in increasing running step rate.<sup>7</sup> Predictably, most of these muscles performed positive work to advance the trail limb forward, while the rectus femoris and tensor fascia latae performed negative work owing to their biarticular attachments.

With increased step rate, muscle forces were observed to increase during late swing, particularly those of the hamstrings and gluteals. This supports previous findings of increased EMG signal of these same muscles during late swing when step rate is increased.<sup>4</sup> During this phase of the running stride, the hamstrings and gluteus maximus accelerate the hip into extension, while the hamstrings also work to oppose the knee from accelerating into extension.<sup>7</sup> As such, the greater activity and force production from these muscles that occur with increased step rate are likely due to increased inertial loads, with muscles contributing to the more erect lower extremity posture that lessens hip and knee joint loads during stance.<sup>13,17</sup> It should be noted that the peak force and negative work of the hamstrings when running at a higher step rate (110% of preferred) are less than 60% of those present during sprinting ( $7.8 \text{ m}\cdot\text{s}^{-1}$ ),<sup>3</sup> that is, well below the loads associated with hamstring strain injury.<sup>14</sup>



The pattern of negative work preceding positive work for most of the muscles has implications for resistance training prescription. To reflect the energetics of these muscles during running, resistance exercise should involve a similar muscle contraction pattern. A rapid-repetition, stretch-shortening activity of eccentric contraction (negative work), followed immediately by concentric contraction (positive work), is recommended. This type of contraction pattern is clearly evident in several muscles at various times in the running cycle, including early stance (gluteals and piriformis), pre-swing/early swing (psoas, iliacus, and adductor longus and brevis), and late swing (hamstrings). Examples of exercises that may be well suited to reflect the energetics of these muscles during running include A skips (hip flexors),<sup>14</sup> B skips (hip flexor and hamstrings),<sup>14</sup> and split-squat jumps (gluteals).<sup>28</sup> Although the intensity of these exercises can be scaled to the individual through speed of movement, injured runners may require less intense forms of exercise depending on symptom severity and provocation.

Based on the increased swing-phase forces and powers produced by some muscles at a higher running step rate, these same exercises may be recommended as part of the initial gait-retraining process. In particular, resistance training of the hamstrings and several hip flexor muscles may be beneficial to facilitate the

desired step rate, due to greater force requirements at specific phases of the running stride. Despite the increased forces for these muscles being relatively small (less than  $1.5 \text{ N}\cdot\text{kg}^{-1}$ ), each muscle must develop these forces during each step; therefore, high-repetition resistance training may be most appropriate.

Despite not being a defined objective of our study, we observed regional differences in force production within each gluteal muscle, exemplifying the complexity that exists in muscle recruitment under triaxial joint loading conditions (FIGURE 3). In the gluteus maximus, the middle fibers produced the greatest forces during loading response, whereas in the gluteus medius peak forces occurred in the anterior fibers. Peak forces in the gluteus minimus occurred in the posterior fibers. This regional variation in force requirements is likely due to the functional demands of the muscle. For example, during loading response, the gluteus medius primarily functions as a hip abductor to provide vertical support to the body, to which the anterior fibers have the greatest moment arm.<sup>20</sup> Also of note, the anterior fibers of the gluteus medius produced the greatest force of any gluteal muscle, which may partially explain why musculotendon tears within the gluteus medius more commonly occur to the anterior fibers.<sup>5</sup> Further, a regional variation in force production is consistent with experimentally obtained

EMG data of the gluteus minimus during walking, suggesting that the hip-stabilizing role of the fibers may differ.<sup>26</sup> Though we acknowledge that the model employed may not fully represent the underlying details that likely contribute to regional force variations within muscle, we nonetheless find it intriguing that our observations are consistent with EMG findings and common injury location.

While characterizing individual muscle forces and powers during running provides useful insights into muscle function and potential injury risk factors, it is important for us to recognize the limitations of this work. We acknowledge that the findings were based on a generic musculoskeletal model that did not take into account any participant-specific information on muscle strengths or geometries. We also assumed a simple scalar relationship between muscle activation and force, which does not account for muscle force-length and force-velocity effects. Muscle activations at any time step of the running simulations were estimated using numerical optimization. Specifically, a set of activations were found that generated the measured hip, knee, and ankle joint accelerations, while minimizing a sum of muscle-volume weighted squared activations.<sup>12</sup> We previously compared the predicted muscle activation patterns to the EMG recordings of several major lower extremity muscles and found good agreement in bursts and phasing.<sup>17</sup> It should be noted that EMG recordings of some muscles (eg, piriformis and adductor brevis) were not available for comparison. Our findings are based on running at preferred speed, ranging from 2.4 to 3.8 m·s<sup>-1</sup>, and therefore may not be generalizable to faster speeds. Despite temporal patterns of force production being similar, muscle force magnitude does not scale proportionally to speed.<sup>7</sup> Some hip muscles were not included in the model (eg, obturator internus and externus) and others were simplified (eg, gemellus superior and inferior were simplified into a single musculotendon unit), which might have had greater influence on the distri-

bution of muscle forces needed to equilibrate hip rotational moments. Finally, all participants were healthy, experienced runners. While it is pertinent to hypothesize about injury, it is unclear whether all of the results apply to injured populations. Future studies should explore muscle forces in those with injury, including patellofemoral pain, iliotibial band syndrome, and gluteal injuries.

## CONCLUSION

**O**UR FINDINGS PROVIDE UNIQUE INSIGHTS into the biomechanical demands placed on the individual hip muscles during running. Specifically, the peak force produced by the gluteus medius was substantially greater than that of any other hip muscle, including the gluteus maximus. Increasing running step rate led to an increase in hip flexor, hamstring, and hip extensor loading in swing, but, conversely, a substantial decrease in peak force and work during loading response in several primary hip muscles was observed. These results may enable clinicians to support and refine current treatment strategies, including exercise prescription and gait retraining, for running-related injuries. ●

## KEY POINTS

**FINDINGS:** During running, the greatest peak force was produced by the gluteus medius during stance phase and the iliacus during swing phase. In general, running with an increased step rate caused a reduction in peak forces of several muscles during stance and an increase during swing.

**IMPLICATIONS:** These findings provide a more complete description of hip muscle demands during running, which is important for scientifically assessing how specific exercises and gait-retraining strategies may be most effective in injury prevention and recovery.

**CAUTION:** Running mechanics are reflective of healthy individuals at preferred speed and may not be generalizable to injured populations.

**ACKNOWLEDGEMENTS:** *The authors would like to thank Christa Wille for her assistance with data collection, as well as the University of Wisconsin Medical Scientist Training Program and the University of Wisconsin Institute for Clinical and Translational Research.*

## REFERENCES

1. Arnold EM, Ward SR, Lieber RL, Delp SL. A model of the lower limb for analysis of human movement. *Ann Biomed Eng*. 2010;38:269-279. <http://dx.doi.org/10.1007/s10439-009-9852-5>
2. Boyajian-O'Neill LA, McClain RL, Coleman MK, Thomas PP. Diagnosis and management of piriformis syndrome: an osteopathic approach. *J Am Osteopath Assoc*. 2008;108:657-664.
3. Chumanov ES, Heiderscheit BC, Thelen DG. Hamstring musculotendon dynamics during stance and swing phases of high-speed running. *Med Sci Sports Exerc*. 2011;43:525-532. <http://dx.doi.org/10.1249/MSS.0b013e3181f23fe8>
4. Chumanov ES, Wille CM, Michalski MP, Heiderscheit BC. Changes in muscle activation patterns when running step rate is increased. *Gait Posture*. 2012;36:231-235. <http://dx.doi.org/10.1016/j.gaitpost.2012.02.023>
5. Connell DA, Bass C, Sykes CA, Young D, Edwards E. Sonographic evaluation of gluteus medius and minimus tendinopathy. *Eur Radiol*. 2003;13:1339-1347. <http://dx.doi.org/10.1007/s00330-002-1740-4>
6. Delp SL, Hess WE, Hungerford DS, Jones LC. Variation of rotation moment arms with hip flexion. *J Biomech*. 1999;32:493-501.
7. Dorn TW, Schache AG, Pandy MG. Muscular strategy shift in human running: dependence of running speed on hip and ankle muscle performance. *J Exp Biol*. 2012;215:1944-1956. <http://dx.doi.org/10.1242/jeb.064527>
8. Ferber R, Davis IM, Williams DS, 3rd. Gender differences in lower extremity mechanics during running. *Clin Biomech (Bristol, Avon)*. 2003;18:350-357.
9. Fredericson M, Cookingham CL, Chaudhari AM, Dowdell BC, Oestreicher N, Sahrmann SA. Hip abductor weakness in distance runners with iliotibial band syndrome. *Clin J Sport Med*. 2000;10:169-175.
10. Geraci MC, Jr., Brown W. Evidence-based treatment of hip and pelvic injuries in runners. *Phys Med Rehabil Clin N Am*. 2005;16:711-747. <http://dx.doi.org/10.1016/j.pmr.2005.02.004>
11. Hamner SR, Seth A, Delp SL. Muscle contributions to propulsion and support during running. *J Biomech*. 2010;43:2709-2716. <http://dx.doi.org/10.1016/j.jbiomech.2010.06.025>
12. Happee R. Inverse dynamic optimization including muscular dynamics, a new simulation method applied to goal directed movements. *J Biomech*. 1994;27:953-960. [http://dx.doi.org/10.1016/0021-9290\(94\)90267-4](http://dx.doi.org/10.1016/0021-9290(94)90267-4)



# [ RESEARCH REPORT ]

13. Heiderscheit BC, Chumanov ES, Michalski MP, Wille CM, Ryan MB. Effects of step rate manipulation on joint mechanics during running. *Med Sci Sports Exerc.* 2011;43:296-302. <http://dx.doi.org/10.1249/MSS.0b013e3181e3bdf4>
14. Heiderscheit BC, Sherry MA, Silder A, Chumanov ES, Thelen DG. Hamstring strain injuries: recommendations for diagnosis, rehabilitation, and injury prevention. *J Orthop Sports Phys Ther.* 2010;40:67-81. <http://dx.doi.org/10.2519/jospt.2010.3047>
15. Ireland ML, Willson JD, Ballantyne BT, Davis IM. Hip strength in females with and without patellofemoral pain. *J Orthop Sports Phys Ther.* 2003;33:671-676. <http://dx.doi.org/10.2519/jospt.2003.33.11.671>
16. Kim JE, Kim KH. Piriformis syndrome after percutaneous endoscopic lumbar discectomy via the posterolateral approach. *Eur Spine J.* 2011;20:1663-1668. <http://dx.doi.org/10.1007/s00586-011-1764-z>
17. Lenhart RL, Thelen DG, Wille CM, Chumanov ES, Heiderscheit BC. Increasing running step rate reduces patellofemoral joint forces. *Med Sci Sports Exerc.* 2014;46:557-564. <http://dx.doi.org/10.1249/MSS.0b013e3182a78c3a>
18. Lu TW, O'Connor JJ. Bone position estimation from skin marker co-ordinates using global optimisation with joint constraints. *J Biomech.* 1999;32:129-134.
19. Mascal CL, Landel R, Powers C. Management of patellofemoral pain targeting hip, pelvis, and trunk muscle function: 2 case reports. *J Orthop Sports Phys Ther.* 2003;33:647-660. <http://dx.doi.org/10.2519/jospt.2003.33.11.647>
20. Neumann DA. Kinesiology of the hip: a focus on muscular actions. *J Orthop Sports Phys Ther.* 2010;40:82-94. <http://dx.doi.org/10.2519/jospt.2010.3025>
21. Niemuth PE, Johnson RJ, Myers MJ, Thieman TJ. Hip muscle weakness and overuse injuries in recreational runners. *Clin J Sport Med.* 2005;15:14-21.
22. Noehren B, Scholz J, Davis I. The effect of real-time gait retraining on hip kinematics, pain and function in subjects with patellofemoral pain syndrome. *Br J Sports Med.* 2011;45:691-696. <http://dx.doi.org/10.1136/bjsm.2009.069112>
23. Paluska SA. An overview of hip injuries in running. *Sports Med.* 2005;35:991-1014. <http://dx.doi.org/10.2165/00007256-200535110-00005>
24. Piazza SJ, Erdemir A, Okita N, Cavanagh PR. Assessment of the functional method of hip joint center location subject to reduced range of hip motion. *J Biomech.* 2004;37:349-356. [http://dx.doi.org/10.1016/S0021-9290\(03\)00288-4](http://dx.doi.org/10.1016/S0021-9290(03)00288-4)
25. Schache AG, Blanch PD, Dorn TW, Brown NA, Rosemond D, Pandy MG. Effect of running speed on lower limb joint kinetics. *Med Sci Sports Exerc.* 2011;43:1260-1271. <http://dx.doi.org/10.1249/MSS.0b013e3182084929>
26. Semciw AI, Green RA, Murley GS, Pizzari T. Gluteus minimus: an intramuscular EMG investigation of anterior and posterior segments during gait. *Gait Posture.* 2014;39:822-826. <http://dx.doi.org/10.1016/j.gaitpost.2013.11.008>
27. Souza RB, Powers CM. Differences in hip kinematics, muscle strength, and muscle activation between subjects with and without patellofemoral pain. *J Orthop Sports Phys Ther.* 2009;39:12-19. <http://dx.doi.org/10.2519/jospt.2009.2885>
28. Struminger AH, Lewek MD, Goto S, Hibberd E, Blackburn JT. Comparison of gluteal and hamstring activation during five commonly used plyometric exercises. *Clin Biomech (Bristol, Avon).* 2013;28:783-789. <http://dx.doi.org/10.1016/j.clinbiomech.2013.06.010>
29. Taunton JE, Ryan MB, Clement DB, McKenzie DC, Lloyd-Smith DR, Zumbo BD. A retrospective case-control analysis of 2002 running injuries. *Br J Sports Med.* 2002;36:95-101.
30. Tonley JC, Yun SM, Kochevar RJ, Dye JA, Farrokhi S, Powers CM. Treatment of an individual with piriformis syndrome focusing on hip muscle strengthening and movement reeducation: a case report. *J Orthop Sports Phys Ther.* 2010;40:103-111. <http://dx.doi.org/10.2519/jospt.2010.3108>
31. van der Worp MP, van der Horst N, de Wijer A, Backx FJ, Nijhuis-van der Sanden MW. Iliotibial band syndrome in runners: a systematic review. *Sports Med.* 2012;42:969-992. <http://dx.doi.org/10.2165/11635400-000000000-00000>
32. van Gent RN, Siem D, van Middelkoop M, van Os AG, Bierma-Zeinstra SM, Koes BW. Incidence and determinants of lower extremity running injuries in long distance runners: a systematic review. *Br J Sports Med.* 2007;41:469-480; discussion 480. <http://dx.doi.org/10.1136/bjsm.2006.033548>
33. Woltring HJ. A Fortran package for generalized, cross-validatory spline smoothing and differentiation. *Adv Eng Software.* 1986;8:104-113. [http://dx.doi.org/10.1016/0141-1195\(86\)90098-7](http://dx.doi.org/10.1016/0141-1195(86)90098-7)



**MORE INFORMATION**  
[WWW.JOSPT.ORG](http://WWW.JOSPT.ORG)

## GO GREEN By Opting Out of the Print Journal

*JOSPT* subscribers and APTA members of the Orthopaedic and Sports Physical Therapy Sections can **help the environment by "opting out"** of receiving *JOSPT* in print each month as follows. If you are:

- **A *JOSPT* subscriber:** Email your request to [jospt@jospt.org](mailto:jospt@jospt.org) or call the *JOSPT* office toll-free at **1-877-766-3450** and provide your name and subscriber number.
- **APTA Orthopaedic or Sports Section member:** Go to <http://www.apta.org/>, log in, and select **My Profile**. Next click on **Email Management/GoGreen**. Toward the bottom of the list, you will find the **Publications** options and may opt out of receiving the print *JOSPT*. **Please save this preference.**

Subscribers and members alike will continue to have access to *JOSPT* online and can retrieve current and archived issues anytime and anywhere you have Internet access.

TABLE 1

PEAK MUSCLE FORCES DURING RUNNING AT 90%,  
100%, AND 110% OF PREFERRED STEP RATE\*

Muscle/Period	90%	100%	110%	Significant P Value		
				90% Versus 100%	90% Versus 110%	100% Versus 110%
Biceps femoris long head						
Stance	3.65 ± 1.03	3.83 ± 0.98	4.07 ± 0.97	...	.0017	...
Late swing	4.69 ± 0.91	5.19 ± 0.94	5.25 ± 0.90	.0002	.0001	...
Semimembranosus						
Stance	7.95 ± 2.22	8.38 ± 2.04	8.95 ± 2.08	...	.0028	...
Late swing	12.12 ± 2.33	13.54 ± 2.35	13.98 ± 2.23	.0001	.0001	...
Gluteus maximus						
Stance	18.01 ± 3.22	15.80 ± 2.89	14.57 ± 2.86	.0001	.0001	.0093
Late swing	5.41 ± 1.49	5.92 ± 1.60	6.04 ± 1.62	.0045	.0005	...
Gluteus medius						
Stance	34.60 ± 5.23	32.05 ± 4.03	28.65 ± 3.63	.0001	.0001	.0001
Late swing	5.84 ± 1.68	6.71 ± 2.17	7.40 ± 2.25	.0006	.0001	.0060
Gluteus minimus						
Stance	24.25 ± 4.22	22.97 ± 3.65	20.82 ± 3.30	.0017	.0001	.0001
Early swing	8.54 ± 2.84	9.76 ± 2.99	9.52 ± 2.85	.0014	.0116	...
Late swing	3.36 ± 1.10	3.86 ± 1.34	4.25 ± 1.43	.0051	.0001	.0413
Tensor fascia latae						
Early swing	1.82 ± 0.42	1.97 ± 0.38	2.12 ± 0.45	.0045	.0001	.0025
Rectus femoris						
Stance	9.98 ± 2.94	8.93 ± 2.78	7.88 ± 2.84	.0001	.0001	.0001
Early swing	7.28 ± 1.59	8.00 ± 1.43	9.11 ± 1.79	.0002	.0001	.0001
Sartorius						
Early swing	0.38 ± 0.09	0.41 ± 0.08	0.46 ± 0.10	.0088	.0001	.0011
Psoas						
Early swing	12.78 ± 3.63	13.19 ± 2.94	13.36 ± 3.17	...	...	...
Iliacus						
Early swing	17.63 ± 3.96	19.29 ± 3.53	20.43 ± 4.16	.0004	.0001	.0164

Table continues on page A2.

TABLE 1

PEAK MUSCLE FORCES DURING RUNNING AT 90%,  
100%, AND 110% OF PREFERRED STEP RATE\* (CONTINUED)

Muscle/Period	90%	100%	110%	Significant P Value		
				90% Versus 100%	90% Versus 110%	100% Versus 110%
Adductor magnus						
Stance	3.01 ± 0.97	2.69 ± 0.89	2.61 ± 0.97	.0072	.0008	...
Adductor brevis						
Early swing	3.36 ± 0.86	3.62 ± 0.75	3.76 ± 0.76	.0054	.0001	...
Adductor longus						
Early swing	1.57 ± 0.41	1.59 ± 0.36	1.63 ± 0.37	...	...	...
Piriformis						
Stance	3.47 ± 1.13	2.97 ± 0.98	2.56 ± 0.82	.0002	.0001	.0009
Early swing	1.27 ± 0.42	1.53 ± 0.39	1.78 ± 0.37	.0001	.0001	.0001
Late swing	1.20 ± 0.61	1.29 ± 0.57	1.34 ± 0.49	...	...	...
Pectineus						
Early swing	0.69 ± 0.16	0.75 ± 0.14	0.78 ± 0.15	.0004	.0001	...
Semitendinosus						
Stance	0.85 ± 0.25	0.88 ± 0.22	0.94 ± 0.23	...	.0103	...
Late swing	1.34 ± 0.26	1.49 ± 0.26	1.53 ± 0.24	.0001	.0001	...
Gracilis						
Early swing	0.43 ± 0.11	0.45 ± 0.12	0.44 ± 0.12	...	...	...
Late swing	0.28 ± 0.07	0.30 ± 0.07	0.30 ± 0.06	...	.0448	...
Quadratus femoris						
Stance	1.46 ± 0.96	1.17 ± 0.79	1.06 ± 0.69	.0143	.0005	...
Late swing	2.37 ± 0.78	2.34 ± 0.84	2.20 ± 0.69	...	...	...
Gemelli						
Stance	1.81 ± 0.98	1.44 ± 0.80	1.16 ± 0.67	.0012	.0001	.0160
Early swing	2.12 ± 0.56	2.31 ± 0.53	2.65 ± 0.60	.0133	.0001	.0001
*Values are mean ± SD N·kg <sup>-1</sup> unless otherwise indicated. For muscles with multiple peaks, the range queried for each peak is listed based on the period of the running cycle.						

TABLE 2

POSITIVE AND NEGATIVE WORK PERFORMED BY EACH  
MUSCLE ACROSS THE RUNNING CYCLE AT 3 DIFFERENT STEP-RATE  
CONDITIONS (90%, 100%, AND 110% OF PREFERRED)\*

Muscle/Work	90%	100%	110%	Significant P Value		
				90% Versus 100%	90% Versus 110%	100% Versus 110%
Biceps femoris long head						
Positive	0.044 ± 0.017	0.045 ± 0.015	0.045 ± 0.015	...	...	...
Negative	-0.055 ± 0.014	-0.056 ± 0.015	-0.060 ± 0.017	...	.0008	.0029
Semimembranosus						
Positive	0.110 ± 0.038	0.100 ± 0.034	0.100 ± 0.034	...	...	...
Negative	-0.170 ± 0.041	-0.180 ± 0.042	-0.200 ± 0.048	.0478	.0001	.0003
Gluteus maximus						
Positive	0.210 ± 0.066)	0.170 ± 0.058	0.150 ± 0.048	.0003	.0001	...
Negative	-0.051 ± 0.030	-0.033 ± 0.036	-0.022 ± 0.021	.0001	.0001	.0179
Gluteus medius						
Positive	0.260 ± 0.063	0.210 ± 0.048	0.170 ± 0.042	.0001	.0001	.0001
Negative	-0.110 ± 0.038	-0.100 ± 0.033	-0.073 ± 0.030	.0449	.0001	.0001
Gluteus minimus						
Positive	0.150 ± 0.049	0.130 ± 0.044	0.100 ± 0.038	.0001	.0001	.0001
Negative	-0.099 ± 0.033	-0.099 ± 0.029	-0.080 ± 0.025	...	.0001	.0001
Tensor fascia latae						
Positive	0.013 ± 0.005	0.013 ± 0.005	0.012 ± 0.005	...	...	...
Negative	-0.032 ± 0.010	-0.032 ± 0.010	-0.030 ± 0.001	...	.0061	.0085
Rectus femoris						
Positive	0.080 ± 0.029	0.074 ± 0.025	0.067 ± 0.023	.0215	.0001	.0293
Negative	-0.370 ± 0.093	-0.350 ± 0.088	-0.340 ± 0.082	.0007	.0001	...
Sartorius						
Positive	0.015 ± 0.004	0.015 ± 0.003	0.016 ± 0.003	...	...	...
Negative	-0.005 ± 0.002	-0.004 ± 0.002	-0.003 ± 0.001	...	.0001	.0049
Psoas						
Positive	0.099 ± 0.031	0.100 ± 0.024	0.100 ± 0.022	...	...	...
Negative	-0.043 ± 0.019	-0.039 ± 0.015	-0.035 ± 0.016	...	.0026	...
Iliacus						
Positive	0.150 ± 0.033	0.170 ± 0.035	0.190 ± 0.037	.0003	.0001	.0001
Negative	-0.089 ± 0.030	-0.079 ± 0.030	-0.075 ± 0.030	.0071	.0002	...

Table continues on page A4.



TABLE 2

POSITIVE AND NEGATIVE WORK PERFORMED BY EACH  
MUSCLE ACROSS THE RUNNING CYCLE AT 3 DIFFERENT STEP-RATE  
CONDITIONS (90%, 100%, AND 110% OF PREFERRED)\* (CONTINUED)

Muscle/Work	90%	100%	110%	Significant P Value		
				90% Versus 100%	90% Versus 110%	100% Versus 110%
Adductor magnus						
Positive	0.033 ± 0.014	0.030 ± 0.014	0.027 ± 0.014	.0126	.0001	...
Negative	-0.007 ± 0.005	-0.006 ± 0.004	-0.004 ± 0.003	...	.0003	.0044
Adductor brevis						
Positive	0.035 ± 0.014	0.038 ± 0.014	0.037 ± 0.013	...	...	...
Negative	-0.021 ± 0.011	-0.019 ± 0.009	-0.015 ± 0.007	...	.0001	.0071
Adductor longus						
Positive	0.026 ± 0.009	0.028 ± 0.010	0.027 ± 0.009	...	...	...
Negative	-0.015 ± 0.006	-0.013 ± 0.005	-0.011 ± 0.004	.0037	.0001	.0017
Piriformis						
Positive	0.017 ± 0.010	0.013 ± 0.008	0.011 ± 0.006	.0006	.0001	.0200
Negative	-0.013 ± 0.006	-0.011 ± 0.005	-0.010 ± 0.004	.0382	.0002	...
Pectineus						
Positive	0.007 ± 0.002	0.007 ± 0.002	0.007 ± 0.002	.0025	.0004	...
Negative	-0.003 ± 0.002	-0.003 ± 0.001	-0.002 ± 0.001	.0270	.0001	.0496
Semitendinosus						
Positive	0.014 ± 0.005	0.013 ± 0.004	0.013 ± 0.004	...	...	...
Negative	-0.023 ± 0.005	-0.024 ± 0.006	-0.026 ± 0.006	...	.0001	.0005
Gracilis						
Positive	0.012 ± 0.006	0.011 ± 0.006	0.009 ± 0.005	.0037	.0001	.0002
Negative	-0.007 ± 0.002	-0.006 ± 0.002	-0.006 ± 0.002	...	.0070	...
Quadratus femoris						
Positive	0.009 ± 0.007	0.007 ± 0.005	0.007 ± 0.004	.0059	.0008	...
Negative	-0.028 ± 0.012	-0.026 ± 0.011	-0.022 ± 0.010	...	.0003	.0220
Gemelli						
Positive	0.011 ± 0.006	0.009 ± 0.004	0.008 ± 0.003	.0417	.0002	...
Negative	-0.015 ± 0.008	-0.013 ± 0.006	-0.012 ± 0.006	.0085	.0002	...

\*Values are mean ± SD J·kg<sup>-1</sup> unless otherwise indicated.